

LEPL “BATUMI SHOTA RUSTAVELI STATE UNIVERSITY”

Faculty of Natural Sciences and Health Care Department of Biology



Department of Biology

Nana Abashidze

**„Ecobiomorphological and Biochemical Peculiarities of some exotic species of Batumi
Botanical Garden in the Adjara seaside”**

Submitted to get a doctorate degree in biology

Specialty: Plant Biodiversity

Annotation

Scientific Supervisor: F. Chaidze

Academic doctor of biology

Scientific Coordinator: A. Kalandia

Propessor of BSU

Batumi 2021

Introduction

Actuality of the Research; The plant introduction and maintenance of exotic species diversity have always been an important priority of the Batumi Botanical Garden. The exotic species introduction works have been going on in the garden since its establishment. The introduction of such trees and plants in through the subtropical zone of the Black Sea, in particular in Adjara, is associated with the 80s of the nineteenth century. Initiators of such works were: I. Veru, M. D'Alphonse, E. Tatarinov, A. Solovtsev, G. Olinsky, B. Stoyanov, A. Korsakov, the Founder of the Garden I. Klingen and others. A Georgian decorator Jason Gordeziani, who had a higher agronomic education in Europe and later worked as an agronomist-decorator of the Royal Garden of Great Britain, had an important role in the process of creating this garden, Batumi seaside park and ornamental tree-plant nurseries arranged around Batumi at that time. He is the first director of the garden, who came by the invitation of A.Krasnov and gained the worldwide recognition, which was manifested by the erection of his bronze bust in the central part of the Batumi Seaside Park, moreover, one of the central squares of the Batumi Botanical Garden was named after Jason Gordezian.

The study of the ecobiomorphological and biochemical peculiarities of exotic plants, introduced from different continents and countries in the garden, still continues, as one of the scientific-theoretical bases for the introduction of plants, which is an essential part to identify new plant species and introduce them in various fields of agriculture.

The prophylactic measures allow producing of various medicinal-prophylactic products, they strengthen as important function of the exotic plants as the protective function of organ is. So far, this is an important task for this research of observing exotics plants. The ingredients of this protective function include: essential oils, vitamins, dietary fiber, minerals, polyunsaturated fatty acids, amino acids, proteins, peptides, organic acids, phenolic compounds (antioxidants) and others.

Aim of the Research, Purpose and Objective; the aim of the research is to study the ecobiomorphological and biochemical peculiarities of some exotic plants introduced in Batumi

Botanical Garden. As far as, one of the main objectives of the Botanical Gardens is to create collections of the botanical species, characterize them under studies, study bio-ecological peculiarities, name growth and development features, identify possibilities of generative - vegetative propagation and essential oil in studying species, determine comparative characters of the natural distribution areas of the studying species and the soil-climatic conditions of the Adjara seaside;

Scientific Novelty; it must be emphasize that, some growing species in the Batumi Botanical Garden have been studied for the first time. Their bio-ecological and biochemical peculiarities have been analyzed and studied according to their seasonal dynamic growth and development; also, effective methods of propagation and the content of essential oils have been elaborated.

Objects of the Research Venue and Methodology: based on the above, the aim of the research is to study the ecobiomorphological and biochemical peculiarities of some exotic plants introduced in Batumi Botanical Garden. Also, estimating the results of their adaptation in the Adjara seaside and setting development measures. 2 groups of exotic plants (11 species and two varieties) identified as research objects.

The first group includes 6 types of essential oil - Aromatic Species and 2 varieties:

1. Yellow, Canary- or Golden Sassafras - *Doryphora Sassafras* Endl.
2. Star-Anise - *Illicium verum* Hook. F.
3. Manuka, Broom Tea-Tree - *Leptospermum scoparium* J.R.Forst. & G.Forst.
4. Manuka, Broom Tea-Tree var. *chapmannii*-*Leptospermum scoparium* var. *chapmannii* Dorrien. Smith.
5. Manuka, Broom Tea-Tree var. *nichollsii* -*Leptospermum scoparium* var. *nichollsii* (Dorr. Sm.)Ewart
6. Linder, Chinese Spicebush - *Lindera communis* Hemsl.
7. Magnolia, Banana Shrub - *Magnolia figo* (Lour.) DC.
8. Brazilian peppertree - *Schinus terebinthifolia* Raddii.

The second group includes 5 types of Medicinal and Fruit-Decorative Species

1. Five-leaf Akebia - *Akebia quinata* (Houtt.) Decne.
2. Cudrang, Chinese Mulberry – *Maclura tricuspidata* Carrière
3. Black Cherry - *Prunus serotina subsp. capuli* (Cav. ex Spreng.)McVaugh.
4. Cattley Guava - *Psidium cattleianum* Afzel. ex Sabine.
5. Six-leafed Stauntonia - *Stauntonia hexaphylla* Decne.

To achieve the goal, we set up the following tasks:

- Characterizing and comparing of the natural distribution areas of the research species and the soil-climatic conditions of the Adjara seaside;
- Determining the possibilities of adaption the introduced species to new environmental conditions and analyzing obtained results;
- Studying ecobiomorphological peculiarities of the research objects and determining the specificity of the process of phonological phases;
- Studying generative peculiarities of the research objects;
- Determining content of the essential oil in leaves, fruit, seeds, flowers of the research objects.

Research Methodology; Different methods have been applied for different purposes in this research: the Mayer Photo-climatic analogue method (Mayer, 1909) for determining the degree of adaptation of introductions; I. Serebryakov (Serebryakov; 1974; 92); I. Beideman (Beideman; 1974; 55); N. Nesterovich (Nesterovich; 1950; 130), A.Gursky (Gursky; 1951: 303) methods for studying the rhythm of growth & development, the timing of fruiting and studying of sprouts biology; I. Elagin (Elagin; 1974, 95) method for clarification of the phonological phases of plants; F. Urbach method (Urbach; 1975; 94) for calculating the percentage of sprouting and flourishing; the "Decorative Dendrology" of M.Tkavadze (Tkavadze; 1990; 5-12) for determining the biometric indicators, the dispersion analysis for mathematical-statistical processing of experimental material (V.Dospekhov; 1985; 351) and Biological method (Gerasikin; 2010, 206) for determining the viability of the rowing material (Standard;1954, 42). The Hydro-distillation,

titration, gas chromatography and mass spectrometric methods have been applied for quantitative and qualitative analysis of essential oils in the studied species.

Material-Technical Bases; the phonological observations of objects under this research were conducted on the specimens growing in the open soil plots of the Batumi Botanical Garden. The propagation works were accomplished in the orangery of the exotic plant department of the Garden. According to the purpose of this research and to determine the content of essential oils in the studying species, there have been conducted an experimental research in the regional chromatographic center of western Georgia.

The dissertation is represented on 141 printed pages, consists 3 chapters, 11 sub-chapters, with 17 tables and 48 colored photos, 160 title lists of references. The main text includes: introduction, literature review, analysis of the results, experimental part, conclusion and recommendations and the list of literature references. The bibliography consists of 160 works by domestic and foreign authors, of which 36 are Georgian, 124 are in foreign languages.

Literature Review; the first chapters of the dissertation analyzes the results of literature and refers the material research. There are also, some discussions about characterization of the soil-climatic conditions of the Batumi Botanical Garden and the area of natural habitat of the studying plants; the regularities of plant introduction, the list of methods used during the studying period; the dissertation is supplied with the list of the literature reference as well.

Chapter I Results Analysis and Outcome of Studying, Botanical-Morphological Characterization of the Exotic Species And Dendrometric Indicators

2 species out of 13 exotics plants growing in the Batumi Botanical Garden have been chosen for the research purpose, they are referred as following; evergreen lianas: Akebia Quinata and Stauntonia Hexaphylla. Also, 7 species: Doryphora Sassafras, Illicium Verum, two varieties of Leptospermum Scoparium (Leptospermum Scosparium var. Chapmannii,

Leopospermium var.); Nichollsii, Psidium Cattleianum, Schinus Terebinthifolia, which are evergreen woody exotics. And, two species: Cudrania Tricuspidata, Prunus Serotina subsp. Capuli, which is a deciduous woody exotic plant. The Latin names of research objects, life form, status, and natural distribution area are given in Table №1 and Dendrometric parameters, year of introduction are given in Table № 2.

	Species	Family	Year of Introduction	Age	Quantity	Height/M/Average/	Stem Diameter- on 1,3 M./cm/Average/	Canopy Circumference/ M/Average/
1	<i>Akebia quinata</i>	<i>Lardizabalaceae</i>	1913	106	2		6 x 7	
2	<i>Maclura tricuspidata</i>	<i>Moraceae</i>	1937	82	15	4	16 x 20	5x3
3	<i>Doryphora sassafras</i>	<i>Atherospermataceaea</i>	1978	42	1	4,5	7 x 9	4 x 3,20
4	<i>Illicium verum</i>	<i>Schisandraceae</i>	1959	61	1	4	12 x 13	3 x 2,80
5	<i>Leptospermum scoparium</i>	<i>Myrtaceaea</i>	1912	107	35	4, 50	13x14	3,75 x 2,80
6	<i>Leptospermum scoparium</i> var. <i>chapmannii</i>	<i>Myrtaceaea</i>	1912	107	25	5, 15	11x 13	5,40 x 3, 46
7	<i>Leptospermum scoparium</i> var. <i>nichollsii</i>	<i>Myrtaceaea</i>	1912	107	20	3	21x 20	3,7 x 2,6
8	<i>Lindera communis</i>	<i>Lauraceae</i>	1958	61	3	3,7	7x23	5,08 x 7,75
9	<i>Magnolia figo</i>	<i>Magnoliaceae</i>	1903	116	17	5	10 x 9	9,0 x 1 ,00
10	<i>Prunus serotina</i> subsp. <i>capuli</i>	<i>Rosaceae</i>	1972	47	5	5	21x 29	8 x 3
11	<i>Psidium cattleianum</i>	<i>Myrtaceaea</i>	1978	41	40	2	35,0 x 40	2 x 3
12	<i>Schinus terebinthifolia</i>	<i>Anacardiaceae</i>	1973	46	1	7	50 x 55	7,3 x 8,6
13	<i>Stauntonia hexaphylla</i>	<i>Lardizabalaceae</i>	1913	106	1		5 x 7	-----

Table 1. Status, Life form and Origin of the of Research Species

Species	Family	Origin	Life Form	Use
<i>Akebia quinata</i> (Houtt.) Decne.	Lardizabalaceae	Japan, Korea, China	Evergreen Liana	Healing, food
<i>Doriphora sassafras</i> Endl.	Atherospermataceae	New South Wales	Evergreen Tree	Healing
<i>Illicium verum</i> Hook. F.	Schisandraceae	China	Evergreen Tree	Healing
<i>Leptospermum scoparium</i> J.R.Forst&G. Forst.)	Myrtaceae	New Zealand	Evergreen tree or Shrub	Healing.
<i>Leptospermum scoparium</i> v. <i>chapmannii</i> Dorien. (Smith.)	Myrtaceae	New Zealand	Evergreen tree or Shrub	Healing.
<i>Leptospermum scoparium</i> v. <i>nicholisii</i> (Dorr.Sm.)Ewart	Myrtaceae	New Zealand	Evergreen tree or Shrub	Healing.
<i>Lindera communis</i> Hemsl.	Lauraceae	South China	Evergreen tree or Shrub	Healing,
<i>Maclura tricuspidata</i> Carrière	Moraceae	China, Himalaya, India	Deciduous Tree	Healing, food
<i>Magnolia figo</i> (Lour.) DC.	Magnoliaceae	South China	Evergreen Tree	Healing
<i>Prunus serotina</i> subsp. <i>capuli</i> McVaugh.	Rosaceae	Peru, Venezuela	Deciduous Tree	Food
<i>Psidium cattleianum</i> Afzel.ex Sabine	Myrtaceae	South America	Evergreen tree or Shrub	Food
<i>Schinus terebinthifolia</i> Raddii.	Anacardiaceae	Brazil, Argentina	Evergreen Tree	Food, Healing
<i>Stauntonia hexaphylla</i> Decne.	Lardizabalaceae	Southern Japan, Korea	Evergreen Liana	Food, Healing

Table 2. Dendrometric Data Obtained through Observations.

Throughout the research, there is given detailed description of botanical-morphological and general biological characterization of each species in their natural habitats (based on reference literature) and in the Batumi Botanical Garden (based on the direct observations and descriptions). Here, in Table №2, there is provided dendrometric data obtained through observations.

Growth and Development Peculiarities of Exotic Species

Studying of the growth and development peculiarities of exotic species, especially introduced species related to the climatic conditions, has an utmost significance, inasmuch as, the seasonal development of plants is an integral indicator of their adaptation. We have been studied the growth and development peculiarities of 13 species in relation to the climatic conditions for five years (2015-2020) on the basis of phonological observation.

The process of vegetative-generative development observations have been conducted through the study of each phonological phase. During the vegetative development: swelling and spreading of vegetative buds; beginning and end of sprout growth (I, II ...); sprout wooding; germination, opening/spreading, reaching final size and massive leaving of young leaves (needles); color change of leaves (needles), leaf falling. During the generative development: swelling and spreading of generative buds; beginning of flowering, massive flowering, beginning and end of pollen dispensation; knotting of fruit (cone), reaching final size, full ripening; pollen dispensation terms. There has been calculated the size of annual growth for each species. The dynamics of leaves and apical growth is given according to decades. During the intensive growth periods the observations have been conducted according to decades, whereas during the less intensive periods which might be considered as twice a month. The great importance had been given to the fruiting process, as much as the intensive fruiting is the main indicative factor of the adaptation degree of the introduced plants in the new environment. The growth and vegetative development of studying species, flowering and fruiting are given in the tables of dissertation, which show peculiarities and terms of each phenol-phase process in 2015-2020 years.

They consistently go through all the stages of vegetative and generative development, some of them even have a self-renewal ability (*Akebia quinata*, *Stauntonia hexaphylla*, *Maclura tricuspidata*), which is an indicator of successful adaptation.

According to the duration of shoot/sprout growth there can be distinguished 3 groups:

Group I - Species that are abundantly fruitful and give seed or sprout and have two periods of the sprout growth: *Doryphora sassafras*, *Maclura tricuspidata*, *Magnolia figo* and *Psidium cattleianum* - 162 days. The first growth of *Doryphora sassafras* shoot starts in spring. The second shoot starts in the second decade of summer.

The first growth of *Maclura tricuspidata* shoot starts in spring and covers 89 day growth period. The second shoot starts in the second decade of summer, covers 65 day growth period and finishes in first decade of autumn. The first growth of *Magnolia figo* shoot starts in the first decade of April and finishes in the second decade of July covers 90 day growth period, while the secondary growth of the sprout begins in the first decade of July and finish in the second decade of August covers 34 day growth period. The first growth of *Psidium cattleianum* shoot starts in spring, covers 70 days growth period and finishes in second decade of July. The second shoot starts in the third decade of August, covers 66 days and finishes in the second decade of autumn.

Group II- includes species with three periods of the sprout growth. The first growth of *Lindera communis* shoots starts in the second decade of April and lasts until the second decade of May. The second growth starts, after a short break, in the second decade of July and lasts until the second decade of August. The third period of growth starts, after a short break, from the second decade of September and finishes about after two months. The vegetative period of the *Lindera communis* is 128 days.

The species in the **group III** normally are the ones with long period of the sprout growth: *Akebia quinata* - duration of shoot growth is 161 days, *Illicium verum*-110days, *Leptospermum scoparium*-170 days, *Leptospermum scoparium* var. *chapmannii* -170 days, *Leptospermum scoparium* var. *nichollsiii* -173 days, *Prunus serotina* subsp. *capuli* - 155 days, *Schinus terebinthifolia* - 255 days, *Stauntonia hexaphylla* - 204 days.

According to the flowering period there can be distinguished **5 groups**: (Table №3)

1. Winter and early spring flowering: *Doryphora sassafras*.

2. Spring flowering: *Akebia quinata*, *Lindera communis*, *Magnolia figo*, *Prunus serotina* subsp. *capuli*.
3. Late spring, summer and Autumn flowering: *Leptospermum Scoparium*, *Leptospermum Scoparium* var. *Chapmannii*, *Leptospermum Scoparium* var. *nichollsii*, *Maclura tricuspidata*.
4. Summer flowering: *Psidium cattleianum*, *Schinus terebinthifolia*.
5. Summer and Autumn flowering: *Illicium verum*.

№	Species	Flower buds		Flowering			Flowering Duration	Flowering Quality
		Swell	Bloom	Start	Massive	finish		
1	<i>Akebia quinata</i>	25,03	12, 04	12, 04	21, 04	10, 05	31	Medium
2	<i>Doryphora sassafras</i>	18,1	7,01	7,01	26, 02	13,04	96	Medium
3	<i>Illicium verum</i>	15,05	15,06	12,06	15,07	18, 12	184	High
4	<i>Leptospermum scoparium</i>	15 , 04	2 ,05	5,10	10 ,06	7, 07	131	High
5	<i>Leptospermum scoparium</i> var. <i>chapmannii</i>	15 , 04	2,05	9 ,05	13, 06	15,07	131	High
6	<i>Leptospermum scoparium</i> var. <i>nichollsii</i>	15, 04	6,05	6,05	11, 06	12,07	131	High
7	<i>Lindera communis</i>	5,03	6,04	7, 04	21,04	15,05	39	High
8	<i>Maclura tricuspidata</i>	11, 03	10,05	14, 05	3, 06	14,09	124	High
9	<i>Magnolia figo</i>	7,03	20, 04	20, 04	24, 05	7,06	49	High
10	<i>Prunus serotina</i> subsp. <i>capuli</i>	8,03	19, 03	16, 04	17, 05	29,05	44	High
11	<i>Psidium cattleianum</i>	12, 04	4,06	5,06	25,06	12,07	38	High
12	<i>Schinus terebinthi folia</i>	5,04	20,04	21,05	15, 06	5,07	46	High
13	<i>Stauntonia hexaphylla</i>	7,03	20,03	29,03	20,04	5,05	43	Medium

Table 3. Flowering of the Observed Species in 2015-2020

The fruiting peculiarities of observed species in new environmental conditions are the integral indicator of this research. There have been singled out 3 groups while evaluating the fruiting process of observed species under the conditions in the Batumi Botanical Garden.

Group I- species with sustained (abundant) fruiting; they are: *Illicium verum*, *Leptospermum scoparium*, *Leptospermum scoparium* var. *chapmannii*, *Leptospermum*

Scoparium var. *nichollsii*, *Maclura tricuspidata*, *Psidium cattleianum*, *Schinus terebinthifolia*, *Stauntonia hexaphylla*.

Group II-species with medium fruiting; they are: *Magnolia figo*, *Doryphora sassafras* *Prunus serotina* subsp. *capuli*.

group III-species with poor fruiting; they are: *Akebia quinata*.

The amplitude reaches the significant value between the limitations of fruiting in some species: *Akebia quinata* (54 days), *Maclura tricuspidata* (64 days), *Doryphora sassafras* (37 days), *Lindera communis* (73 days), and *Psidium cattleianum* (39 days), *Schinu terebinthipholia* (57 days), *Stauntonia hexaphylla* (32 days), *Illicium verum* (33 days).

The amplitude varies slightly between the limitations of fruiting in the following exotics: *Leptospermum scoparium* (21 days), *Leptospermum scoparium* var. *chapmannii* (9 days), *Leptospermum scoparium* var. *nichollsii* (7 days), *Magnolia figo* (20 days), *Prunus serotina* subsp. *capuli* (8 days).

№	Species	Fruiting Quality			Fruiting Degree
		Start	Massive	Finish	
1	<i>Akebia quinata</i>	7, 09	5,1	11,1	Low
2	<i>Doryphora sassafras</i>	1,07	20,07	8,08	Medium
3	<i>Illicium verum</i>	12,07	16,09	20,12	High
4	<i>Leptospermum scoparium</i>	23, 05	13,06	3, 10	High
5	<i>Leptospermum scoparium</i> var. <i>chapmannii</i>	24, 05	13, 06	4, 10	High
6	<i>Leptospermum scoparium</i> var. <i>nichollsii</i>	24, 05	13,06	3, 10	High
7	<i>Lindera communis</i>	19,09	5, 10	26, 10	High
8	<i>Maclura tricuspidata</i>	22,09	5, 11	2, 12	High
9	<i>Magnolia figo</i>	30, 08	19, 09	6, 10	High
10	<i>Prunus serotina</i> subsp. <i>capuli</i>	8, 07	26, 07	9, 08	Medium
11	<i>Psidium cattleianum</i>	18,08	12,1	2, 11	High
12	<i>Schinus terebinthifolia</i>	10, 10	24, 11	11, 12	High
13	<i>Stauntonia hexaphylla</i>	22, 09	25 10	9, 11	Medium

Table 4. Fruiting of Research Species

Chapter III Results of Generative Multiplication of the Observed Species

One of the main objectives for diversity of plant propagation is to elaborate effective methods for their reproduction. Studying the reproduction biology during the introduction process has the essential theoretical and practical value.

There had been studied an ability of generative-reproduction of the plants through seeds and on this purpose sowing was conducted in two variants in this study:

- 1) Sowing of seeds collected in autumn;
- 2) Sowing of seeds taken in autumn, in early spring,

3.1. Propagation through Seeds

Seed was gathered from the collection of the fruiting exotic species. The studies of the biology of seed propagation, germination and of the species under study were accomplished in the orangery.

While determining seed productivity of the research species, we have singled out three groups:

1. Species with high germination coefficient;
2. Species with medium coefficient of germination.
3. Species with low or none coefficient of germination.

Group I - species with high germination coefficient belongs: *Maclura tricuspidata*, *Leptospermum scoparium*, *Leptoapermum scoparium* var. *chapmannii*, *Leptoapermum scoparium* var. *nichollsii*, *Lindera communis*, *Magnolia figo*, *Psidium cattleianum*, *Stauntonia hexaphylla*.

Group II - species with medium germination coefficient belongs: *Akebia quinata* and *Illicium verum*.

Group III - species with low germination coefficient or not at all belongs: *Prunus serotina* subsp. *capuli*, *Schinus terebinthifolia*.

Species	Seed Quantity	Seed Collection Time	Sowing Time	Emergence			Time Required for Emergence	%
				First	Massive	Final	Day	
<i>Maclura tricuspidata</i>	200	2, 11	24,03	5, 04	30, 04	4,05	20	90±3,2
<i>Leptospermum scoparium</i>	200	12, 06	25,1	24, 03	17, 04	8,05	121	78±2,7
<i>Leptospermum scoparium</i> var. <i>chapmannii</i>	200	12, 06	25,1	24, 03	17, 04	10,1	121	70±3,1
<i>Leptospermum scoparium</i> var. <i>nichollsii</i>	200	12, 06	25,1	24, 03	17, 04	12,1	121	70±2,5
<i>Lindera communis</i>	200	13, 07	25,1	13, 03	19, 04	13,1	62	69±3,4
<i>Magnolia figo</i>	200	6, 10	11,03	5, 04	17, 04	12,1	25	70±3,1
<i>Prunus serotina</i> subsp. <i>capuli</i>	200	26 ,07	25,1	24, 03	8,04	7,05	121	30±1,1
<i>Psidium cattleianum</i>	200	14, 10	25,1	24, 03	9, 04	12,1	121	85±2,4
<i>Schinus terebinthifolia</i>	200	22, 11	24,03	9, 04	24.04.	6,05	16	20±1,3
<i>Stauntonia hexaphylla</i>	200	18, 10	25,1	24, 03	17, 04	30	38	80±2,7

Table 5 Dynamics of Seed Emergence of Research Species

Chapter IV Biochemical Peculiarities of Observed Objects

In General, plant introduction and preservation of diversity have always been a significant priority of botanical gardens. The introduction works in the Batumi Botanical Garden are connected with the foundation of the Garden. Palibin, who was the director of the Garden from 1916, is considered as the founder of the observation activities in the Garden. He had supervised the studies on growth & development peculiarities of the introduced plants and tried to find out how useful our climate conditions might be to spread certain plants or use them in agriculture, decorative gardening and other fields of national economy was observed, though these studies.

The study of Eco-physiological and biochemistry peculiarities and features of introduced plants with various geographical origins still continuous at the Botanical Garden, as well as creating the theoretical basics of plant introduction in order to reveal new plant resources and

implement them in different types of farming economy. The scientific studies of plant introduction require a complex integrated approach, which is an understanding of their close relations with the natural environment in the conditions of interrelations between morphological, phylogenetic and bio-ecological factors along with the knowledge of chemical composition. The members of the rose, myrtle, laurel, cypress family are especially rich in aromatic plants [3]. Objects of our study have been the exotics: *Doryphora sassafras* Endl; *Leptospermum Scoparium* Forst; *Leptospermum Scoparium* var. *Chapmannii*; Dorien Smith Carr; *Leptospermum Scoparium* var. *Nichollsii* (Dorr.Sm.); Ewart; *Illicium Verum* Hook.; *Lindera Communis* Thunb; *Magnolia Figo* (Lour.) DC.

Based on the results of growth and development of the introduced plants studies, researchers have been established that they consist all phases of vegetative and generative development, which shows a successful acclimatization rate.

The goal of our study was to reveal the results of the adaptation of the plants, by observing their bio-ecological and biochemical peculiarities and estimate regularities of development under the cultivation.

In order to determine the content of essential oils, there has been conducted experimental research in the regional chromatographic center of western Georgia. The methods of hydro-distillation, high pressure super-fluid extraction (with inert gases) and titration have been applied to isolate of essential oils.

Through the titration method (hydrolysis of essential oils glycosides link in leaves and fruit was conducted with hydrogen chloride action. The free essential oil was titrated with bromine). It was determined that the content of essential oils in *Leptospermum Scoparium* var. *Chapmannii* Dorien. Smith- leaves is $0,1533 \pm 0,005$ %; *Leptospermum Scoparium* J.R.Forst. & G.Forst – leaves is $0,065 \pm 0,008$ %; *Leptospermum Scoparium* v. *Niichollsii* (Dorr. Sm.) Ewart- leaves is $0,065 \pm 0,007$ %; *Doryphora sassafras* Endl- leaves is $0,124\% \pm 0,02$ %; *Illicium Verum* Hook. f. - Fruits is $0,0657 \pm 0,006\%$; *Lindera Communis* Hemsl- fruits- $0,5474 \pm 0,06$ %, *Magnolia Figo* (Lour.)DC- flowers is $1,078 \pm 0,03$. Table 5.

Species Name	Total Essential Oils %		
	Leaves	Fruits	Flowers
<i>Doryphora sassafras</i>	0,124 ± 0,02		
<i>Illicium Verum</i>		0,0657 ± 0,006	
<i>Leptospermum Scoparium</i> v. <i>Chapmannii</i>	0,1537 ± 0,005		
<i>Leptospermum Scoparium</i>	0,065 ± 0,008		
<i>Leptospermum Scoparium</i> v. <i>Nicholssii</i>	0,065 ± 0,007		
<i>Lindera Communis</i>		0, 5474 ± 0,06	
<i>Magnolia Figo</i>			1.078 ± 0,03

Table № 5 The Quantity of essential Oils Obtained by the Titration Method

4.1. Obtaining an Aromatic Complex of *Leptospermum Scoparium*s leaves, *Illicium Verum*s and *Lindera Communis* fruits

The aromatic foliage complex of *Leptospermum Scoparium* leaves, *Illicium Verum* and *Lindera Communis* fruits were obtained by Hydro-distillation. 70 grams of dried leaf (crushed) and fruits together with 200 ml of water were placed in a flask for the distillation process, the distillation process had been carried out using a Clevenger-type apparatus (Pic.1-2) for 1 hours and the condensation in a refrigerator at temperature - 0.0 ° C. The obtained essential oil was extracted with hexane, 0.5 µl of the organic part of which was centrifuged (2 minutes at 1350 revolutions / min) and injected on the chromatograph.



Picture1. Obtained Essential oils by Clevenger Method



Picture2. Research Sample *Leptospermums*

4.1. Essential oils of Fruit of *Leptospermum Scoparium* leaves, *Illicium Verum* and *Lindera Communis* obtained by Gas Chromatography

The temperature mode of the injector was 300 ° C, while the sample in the column was divided into helium flux by a ratio of 1 / 214.3. The chromatography was carried out in four stages at a temperature gradient. In particular, chromatography was started at 50 ° C, which was continued for 5 minute at that temperature, then increased to 200 ° C (second stage) at a rate of 3.0 (° C / min) and was continued for 5 minute. In the third stage, it increased to 320 ° C, at a speed of 30 (° C / min) and continued at that temperature for 20 minute. The whole chromatographic implementation time was 74.0 minutes. The essential oil components separated by chromatography were detected by an allure-ionization detector. The quantitative content of essential oil was determined with an accuracy of 0.01% in percentage according to the peak area.




Figure 1. Essential Oils of *Leptospermum Scoparium* leaves obtained by Chromatogram

Peak Number №	Peak Time	Component	Area,
	(min)	Name	%
1	8,505	a-Thujene	0.955±0.03
2	23,14	Peak	0.643±0.02
3	28,64	Geranyl acetat	0.585±0.02
4	29,95	a-Cubebene	1.710±0.06
5	31,845	a-Copaene	6.717±0.24
6	32,265	B-Elemene	23.238±0.81
7	33,28	Menthyl acetat	3.307±0.12
8	33,59	a-Sellnene	0.855±0.03
9	34,475	Peak	3.296±0.12
10	36,21	Dehydroelsholt	11.556±0.40
11	36,465	Peak	12.321±0.43
12	37,215	Peak	4.214±0.15
13	37,485	Peak	8.568±0.30
14	37,865	Peak	2.273±0.08
15	38,73	Peak	0.816±0.03
16	39,84	Peak 55	3.049±0.11
17	40,06	Peak	3.070±0.11
18	41,675	Peak	1.794±0.06
19	41,88	Peak	0.924±0.03
20	42,22	Peak 58	2.054±0.07
21	42,845	Peak 59	4.749±0.17
22	42,99	Peak	3.307±0.12

Table 6. The Component Composition of the Essential Oil in the Leaves of *Leptospermum Scoparium*

As a result of the chromatographic research, 22 components have been found in the essential oils of *Leptospermum Scoparium* leaves. Among them, there were identified 8 components, In particular: α -Thujene 0.955 ± 0.03 %, Geranyl Acetat 0.585 ± 0.02 %, α -Cubebene 1.710 ± 0.06 %, α -Copaene 6.717 ± 0.24 %, B-Elemene 23.238 ± 0.81 %, Menthyl Acetat 3.307 ± 0.12 %, α -Sellnene 0.855 ± 0.03 %, Dehydroelsholt 11.556 ± 0.40 %.




Figure 2. Chromatogram of *Lindera Communis* fruits Essential oils

Peak №	Peak Time (min)	Component Name	Area %
1	19.220	Peak	0.230 ± 0,0069
2	20.125	26	0.070 ± 0,0021
3	21.545	Peak	0.369 ± 0,01107
4	21.995	27	0.173 ± 0,0051
5	24.122	Perilaldehyde	0.075 ± 0,00225
6	24.425	1, 8-Cineole	4.420 ± 0,1326
7	25.460	Peak	92.017 ± 2,7605
8	26.360	30	0.036 ± 0,0010
9	26.980	Citronelylacet	0.012 ± 0,00036
10	27.250	Peak	0.040 ± 0,0012
11	27.665	Neryl acetate	0.112 ± 0,0034
12	28.130	α-Copaene	0.025 ± 0,0008
13	28.715	Geranyl acetat	0.054 ± 0,0016
14	28.935	Peak	0.084 ± 0,0025
15	29.155	Peak	0.149 ± 0,0046
16	29.470	Peak	0.019 ± 0,0005
17	29.595	Dodecanal	0.023 ± 0,0007
18	30.620	Peak	0.017 ± 0,0005
19	32.645	33	0.020 ± 0,0006
20	33.000	2 (E) -Dodecenal	0.064 ± 0,0019
21	36.455	Peak	0.032 ± 0,0010
22	40.895	Peak 56	0.262 ± 0,0079
23	42.335	Peak 58	0.362 ± 0,0109
24	42.685	α-Isocomene	0.009 ± 0,0003
25	43.860	α-Terpineol ac	0.270 ± 0,0081
26	44.475	Peak 62	0.051 ± 0,0015
27	45.280	Peak	0.547 ± 0,0164
28	45.440	Peak	0.026 ± 0,0008
29	45.530	Peak	0.203 ± 0,0061
30	45.645	Peak	0.037 ± 0,0011
31	49.115	E-Caryophyllen	0.050 ± 0,0015
32	50.210	Peak 46	0.029 ± 0,0009
33	51.220	Peak	0.028 ± 0,0008
35	51.905	3	0.021 ± 0,0006
36	52.220	Peak	0.050 ± 0,0015
37	53.715	4	0.005 ± 0,0002

Table 7. The Component Composition of *Lindera Communis* Fruits Essential Oil

As a result of our chromatographic research, 37 components have been found in the essential oils of *Lindera Communis* fruits. Among them there were identified 11 , In particular:

Perilaldehyde $0.075 \pm 0,00225$ %, 1,8-Cineole $4.420 \pm 0,1326$ %, Citronelylacetat $0.012 \pm 0,00036$ %, Nerylacetate $0.112 \pm 0,0034$ %, α -Copaene $0.025 \pm 0,0008$ %, Geranyl acetate $0.054 \pm 0,0016$ %, Dodecanal $0.023 \pm 0,0007$ %, 2(E)-Dodecenal $0.064 \pm 0,0019$ %, α -Isocomene $0.009 \pm 0,0003$ %, α -Terpineol acetate $0.270 \pm 0,0081$ %, (e)-beta-Caryophyllene $0.050 \pm 0,0015$ %.




Figure 2. Chromatogram of *Illicium Verum* fruits Essential oils

Peak №	Peak Time (min)	Component Name	Area %
1	19.225	Peak	1.838 ± 0,0551
2	21.540	Peak	0.108 ± 0,0032
3	21.990	27	0.645 ± 0,0194
4	24.110	Perilaldehyde	0.941 ± 0,0282
5	24.355	1, 8-Cineole	1.188 ± 0,0356
6	24.885	Peak	7.728 ± 0,2318
7	25.575	Peak	0.216 ± 0,0065
8	26.915	Citronelylacet	0.563 ± 0,0169
9	27.675	Neryl acetate	4.585 ± 0,1375
10	31.805	32	0.821 ± 0,0246
11	32.575	33	4.802 ± 0,1441
12	34.705	Thymol methyl	0.263 ± 0,0079
13	34.875	(E,E)-α-Farnes	0.509 ± 0,0153
14	40.360	Silphinene	0.702 ± 0,0211
15	40.605	Peak	0.247 ± 0,0074
16	40.920	Peak 56	5.861 ± 0,1758
17	42.370	Peak 58	10.076 ± 0,3023
18	43.675	Z-Caryophyllen	0.282 ± 0,0085
19	43.870	α-Terpineol ac	0.852 ± 0,0256
20	44.505	Peak	2.553 ± 0,0766
21	44.750	Peak 62	2.885 ± 0,0865
22	44.860	Peak	0.219 ± 0,0066
23	45.280	Peak	2.169 ± 0,0651
24	45.475	Peak	1.121 ± 0,0336
25	45.530	Peak	1.980 ± 0,0594
26	45.695	Peak	2.221 ± 0,0666
27	46.285	Peak 63	0.938 ± 0,0282
28	46.790	Peak 65	0.059 ± 0,0018
29	47.410	Peak 67	0.370 ± 0,0111
30	47.815	Peak 68	1.253 ± 0,0376
31	48.045	Peak 70	0.134 ± 0,0040
32	48.615	Peak 71	0.312 ± 0,0094
33	48.910	Peak	1.326 ± 0,0398
34	49.130	E-Caryophyllen	0.780 ± 0,0234
35	49.255	Peak	0.939 ± 0,0282
36	49.625	Peak 72	1.192 ± 0,0358
37	50.040	Peak 73	0.520 ± 0,0156
38	50.480	Peak 46	1.053 ± 0,0316
39	50.670	Peak	0.114 ± 0,0034
40	50.750	Peak	0.063 ± 0,0019
41	50.965	Peak	0.996 ± 0,0299
42	51.310	Peak	0.960 ± 0,0288
43	51.530	Peak	4.860 ± 0,1458
44	51.925	3	3.731 ± 0,1119
45	52.135	Peak	5.564 ± 0,1669
46	52.500	Peak 74	18.073 ± 0,5421
47	53.735	4	0.546 ± 0,0164
48	53.850	Peak	0.237 ± 0,0071
49	54.525	Peak 48	0.604 ± 0,0181

Table 8. The component composition of the essential oil in the fruits of *Illicium Verum*

As a result of our chromatographic research, 49 components have been found in the essential oils of *Illicium Verum* fruits. Among them there were identified 10, In particular:

Perilaldehyde $0.941 \pm 0,0282$ %, 1, 8-Cineole $1.188 \pm 0,0356$ %, Citronelylacetat $0.563 \pm 0,0169$ %, Nerylacetate $4.585 \pm 0,1375$ %, Thymol methyl ether $0.263 \pm 0,0079$ %, (e)-alpha-farnesene $0.509 \pm 0,0153$ %, Silphinene $0.702 \pm 0,0211$ %, Z-Caryophyllene $0.282 \pm 0,0085$ %, Alpha-Terpinyl Acetat $0.852 \pm 0,0256$ %, (e)-beta-Caryophyllene $0.780 \pm 0,0234$ %.

Chapter V Conclusions and Recommendations

On the basis of the conducted introductive research, while studying morphological peculiarities of separate species, vegetation period the peculiarities of seasonal growth and development, reproduction issues, terms of separate phenol-phases, period of dormancy, winter resistance, heir aromatic, ornamental, medicinal nutritious and other economic-biological characteristics were evaluated.

1. According to the growth and developmental characteristics of the observed species, it has been concluded that they go through all the stages of vegetative and generative development in sequence, which indicates their successful adaptation. For the purpose of research, there has been carried out phonological observations on the experimental plants. Interrelation between the flowering and fruit bearing was estimated in related to the internal and external factors. Furthermore, plant development phases, air temperature, drought- resistance, seed quality (germination capacity) and other indices were evaluated. The studies of seed productivity have established that the coefficient of seed emergence is quite high. they consistently go through all the stages of vegetative and generative development, some of them even have a self-renewal ability (*Akebia quinata*, *Stauntonia hexaphylla*, *Maclura tricuspidata*), which is an indicator of successful adaptation.

2. According to the duration of sprout growth 3 groups have been singled out;

Group I- species with two period of shoot growth: shoot growth period of *Doryphora sassafras* - 180 days, *Maclura tricuspidata* 114 days *Magnolia Figo* 122 days and *Psidium cattleianum* - 136 days.

Group II - species with three period of sprout growth: *Lindera communis* growth period is 180 days.

Group III -species with a long period of shoot growth: *Akebia quinata* - duration of shoot growth is 161 days, , *Illicium Verum* - 110 days, *Leptospermum scoparium* - 170 days, *Leptospermum scoparium* var. *chapmannii* – 170 days, *Leptospermum scoparium* var. *nichollsii* -173 days, *Prunus serotina* subsp. *capuli*- 152 days, *Schinus terebinthifolia* - 213 days, *Stauntonia hexaphylla* - 233 day.

3. According to the flowering period five groups have been singled out on the basis of observation:

- ***Winter and early spring flowering:*** *Doryphora Sasafras*.
- ***Spring flowering:*** *Akebia Quinata*, *Lindera Communis*, *Magnolia Figo*, and *Prunus Srtotina* subsp. *Capuli*.
- ***Late spring and summer flowering:*** *Leptospermum Scoparium*, *Leptospermum Scoparium* var. *Chapmannii*, *Leptospermum Scoparium* var. *Nichollsii*.
- ***Summer flowering:*** *Psidium Cattleianum*, *Schinus Terebinthifolia*.
- ***Summer and autumn flowering:*** *Illicium Verum*.

4. In assessing the fruiting in new environmental conditions, it is established that the fruiting of introduced species in the conditions of the Batumi Botanical Garden is stable and the quality is quite high. ***According to the degree of fruiting there can be distinguished 3 groups of species:***

Group I -species with sustained (abundant) fruiting: *Maclura tricuspidata*, *Illicium verum*, *Leptospermum scoparium*, *Leptospermum scoparium* var. *chapmannii*, *Leptospermum scoparium* var. *nichollsii*, *Psidium cattleianum*, *Schinus terebinthifolia*, and *Stauntonia hexaphylla*.

Group II- species with medium fruiting: *Magnolia Figo*, *Doryphora Sassafras*, *Prunus serotina* subsp. *capuli*.

Group III -species with poor fruiting: *Akebia quinata*.

The amplitude reaches a significant value in assessing the limitations of fruiting in some species: *Akebia quinata* (54 days), *Maclura tricuspidata* (64 days), *Doryphora sassafras* (37 days), *Lindera communis* (73 days), *Psidium cattleianum* (39 days), *Schinus terebinthifolia* (57 days), *Stauntonia hexaphylla* (32 days), and *Illicium verum* (33 days). The amplitude varies slightly in assessing the limitations of fruiting in the following exotics: *Leptospermum scoparium* (21 days), *Leptospermum Scoparium* var. *Chapmannii* (9 days), *Leptospermum Scoparium* var. *Nichollsii* (7 days), *Magnolia Figo* (20 days), *Prunus Serotina* subsp. *Capuli* (8 days).

5. In According to determine the seed productivity of the observed species *there can be distinguished 3 groups of species:*

1. Species with high germination coefficient;
2. Species with medium coefficient of germination.
3. Species with low or none coefficient of germination.

The first group of species with high germination coefficient belongs: *Maclura tricuspidata*, *Leptospermum scoparium*, *Leptospermum scoparium* var. *chapmannii*, *Leptospermum scoparium* var. *nichollsii*, *Lindera communis*, *Magnolia figo*, *Psidium cattleianum*, and *Stauntonia hexaphylla*.

The second group of species with medium germination coefficient belongs: *Akebia quinata*, *Illicium verum*.

The third group of species with low germination coefficient or none at all belongs: *Prunus serotina* subsp. *capuli*, *Schinus terebinthifolia*.

6. For the purpose of determining the content of essential oils there has conducted experimental research in the regional chromatographic center of western Georgia. For the isolation of essential oils the methods of hydro-distillation, high pressure super-fluid extraction (with inert gases) and titration have been applied. Through the titration method (hydrolysis of essential oils glycosides link in leaves and fruit was conducted with hydrogen chloride action. The free essential oil was titrated with bromine). It was determined that the content of essential oils in the *Leptospermum scoparium* var. *chapmannii* leaves $0,1533 \pm 0,005$ %, *Leptospermum scoparium*. leaves $0,065 \pm 0,008$ %, *Leptospermum scoparium* v. *nichollsii*

leaves $0,065 \pm 0,007$ %, *Doryphora sassafras* leafs $0,124\% \pm 0,02$ %, *Illicium verum* fruits - $0,0657 \pm 0,006\%$, *Lindera communis* fruits $-0,5474 \pm 0,06$ %, *Magnolia figo* flowers $1,078 \pm 0,03$.

7. As a result of our chromatographic research, 22 components have been found in the essential oils of *Leptospermum scoparium* leaves. Among them there were identified 8 components, In particular: α -Thujene 0.955 ± 0.03 %, Geranyl Acetat 0.585 ± 0.02 %, α -Cubebene 1.710 ± 0.06 % , α -Copaene 6.717 ± 0.24 %, B-Elemene 23.238 ± 0.81 %, Menthyl Acetat 3.307 ± 0.12 %, α -Selinene 0.855 ± 0.03 %, Dehydroelsholt 11.556 ± 0.40 %.

8. As a result of our chromatographic research, 37 components have been found in the essential oils of *Lindera communis* fruits. Among them there were identified 11 components, In particular: Perilaldehyde $0.075 \pm 0,00225$ %, 1,8-Cineole $4.420 \pm 0,1326$ %, Citronelylacetat $0.012 \pm 0,00036$ %, Nerylacetate $0.112 \pm 0,0034$ %, α -Copaene $0.025 \pm 0,0008$ %, Geranyl acetate $0.054 \pm 0,0016$ %, Dodecanal $0.023 \pm 0,0007$ %, 2(E)-Dodecenal $0.064 \pm 0,0019$ %, α -Isocomene $0.009 \pm 0,0003$ %, α -Terpineol acetate $0.270 \pm 0,0081\%$, (e)-beta-Caryophyllene $0.050 \pm 0,0015\%$.

9. As a result of our chromatographic research, 49 components have been found in the essential oils of *Illicium verum* fruits. Among them there were identified 10 components, are Perilaldehyde $0.941 \pm 0,0282$ %, 1, 8-Cineole $1.188 \pm 0,0356$ %, Citronelylacetat $0.563 \pm 0,0169$ %, Nerylacetate $4.585 \pm 0,1375$ %, Thymol methyl ether $0.263 \pm 0,0079$ %, (e)- α -farnesene $0.509 \pm 0,0153$ %, Silphinene $0.702 \pm 0,0211\%$, Z-Caryophyllene $0.282 \pm 0,0085$ %, α -Terpinyl Acetat $0.852 \pm 0,0256$ %, (e)-beta-Caryophyllene $0.780 \pm 0,0234$ %.

10 . The some aromatic and medicinal plant species introduced at Batumi Botanical Garden take a significant place in terms of their application in medicine and can widely be implemented in modern pharmacology. According to the adaptation degree, the studied plants can be recommended for the reasonable application in the Adjara coastline.

Chapter VI Recommendations

Some exotic plants introduced in the soil-climatic conditions of Batumi Botanical Garden have high adaptation ability, so we recommend interested legal entities and individuals to distribute for various purposes in the Adjara coast.

Winter flowering decorative exotic plant *Doryphora Sassafras* can be used in ornamental gardening, in medicine, modern pharmacology, as a unit copy or in group plantings, due to the content of essential oils in the leaves. Summer and autumn flowering *Illicium verum* can be widely used in decorative gardening, as far as the fruit contains an essential oil, it also be it can be used in medicine, cooking and cosmetology.

A decorative exotic, *Leptospermum* (*Leptospermum scoparium*) creates an excellent ensemble, especially with its red and pink varieties. Also they contain essential oils and can be used in decorative gardening, medicine, and cosmetology. It is possible to cultivate agricultural plantations as very valuable exotic.

Spring flowering, aromatic *Lindera communis* and *Magnolia figo* can be used in decorative gardening, medicine, cosmetology as the promising essential oil.

Summer flowering *Schinus terebinthifolia* can be used in the food industry, perfumery.

Lardizabalaceae family members: spring flowering *Akebia quinata* and *Stauntonia hexapylla* are precious fruit-decorative exotic plants and can be used in food industry, medicine and decorative gardening.

Maclura tricuspidata is decorative with beautiful, red fruits.; its fruit contains sugar and natural acids, also, large amounts of B, C, PP vitamins, carotene and rutting. It is also rich in glycosides, pectin, resin substances and iron; therefore it can be widely used in medicine.

Summer-flowering, fruit-decorative exotics: *Prunus serotina subsp. capuli* and *Psidium cattleianum* can be cultivated in agricultural plantations.

Reference

1. აბაშიძე1959: აბაშიძე ი. „დენდროლოგია მცენარეთა გეოგრაფიის ელემენტებით“. ნაწილი I. სასოფლო-სამეურნეო ინსტიტუტის გამომცემლობა. თბილისი. გვ. 119, 113,
2. აბაშიძე2014: აბაშიძე ნ., ჩაიძე ფ. " ბათუმის ბოტანიკურ ბაღში ინტროდუცირებული ეგზოტების ნაყოფმსხმოიარობის თავისებურებანი." ბათუმის ბოტანიკური ბაღის მოამბე 35. ბათუმი. ISSN 1987-8621. გვ. 27-30.
3. აბაშიძე 2015: აბაშიძე ნ. „ბათუმის ბოტანიკურ ბაღში ინტროდუცირებული გვარი ლეპტოსპერმუმის ზოგიერთ წარმომადგენლებში ეთერზეთების შემცველობა“. V სექცია. ბათუმის შოთა რუსთაველის სახელმწიფო უნივერსიტეტის 80 წლის იუბილესადმი მიძღვნილი სტუდენტთა საერთაშორისო სამეცნიერო კონფერენცია. ბათუმის შოთა რუსთაველის სახელმწიფო უნივერსიტეტი. ბათუმი. გვ.16.
4. აბაშიძე...2017: აბაშიძე ნ. ჩაიძე ფ. კალანდია ა. ბათუმის ბოტანიკურ ბაღში ინტროდუცირებულ ზოგიერთ მცენარეთა ბიოეკოლოგიური და ბიოქიმიური თავისებურებები. III სამეცნიერო კონფერენცია „ბიომრავალფეროვნება და საქართველო“. საქართველოს ეროვნული ბოტანიკური ბაღი. თბილისი, გვ. 148-151.
5. აბაშიძე...2019: აბაშიძე ნ. ჩაიძე, ფ. კალანდია ა. ჩიქოვანი ი. „ ბათუმის ბოტანიკურ ბაღში ინტროდუცირებული ზოგიერთი არომატული ეგზოტის ბიოეკოლოგიური

თავისებურებები და მათში ეთერზეთების შემცველობა.“

საქართველოს სოფლის მეურნეობის მეცნიერებათა

აკადემიის მოამბე N 2 (42). ISSN 1512- 2743 . თბილისი. გვ. 119-121.

6. აბაშიძე2019 : აბაშიძე ნ. ბათუმის ბოტანიკურ ბაღში ინტროდუცირებული ზოგიერთი ეგზოტის ადაპტაციის თავისებურებები და გამოყენების პერსპექტივები მედიცინაში, ახალგაზრდა მეცნიერთა და სტუდენტთა კონფერენცია თანამედროვე ბიომედიცინის საკითხები. ISBN -978-9941-8-1697-0. ბათუმი. გვ. 4-6.
7. აბაშიძე2020 : აბაშიძე ნ. „ ფსიდიუმის (*Psidium cattleianum* Afzel.ex Sabine) ეკობიომორფოლოგიური თავისებურებები ბათუმის ბოტანიკურ ბაღში“ საქართველოს სოფლის მეურნეობის მეცნიერებათა აკადემიის მოამბე N 2 (44). ISSN 1512-2743. თბილისი. გვ. 83– 85.
8. ბალანჩივაძე... 1961: ბალანჩივაძე შ. „ახალი ზელანდიის მცენარეთა აკლიმატიზაცია საქართველოს შავი ზღვის სანაპიროზე“. სახელმწიფო გამომცემლობა, ბათუმი. გვ. 44—55.
9. ბალანჩივაძე ...1968: შ. ბალანჩივაძე „მექსიკის მერქნიან მცენარეთა აკლიმატიზაციის შედეგები ბათუმის ბოტანიკურ ბაღში“. ბათუმის ბოტანიკური ბაღის მოამბე №13 გამომც. „მეცნიერება“, თბილისი. გვ. 27.
10. ბასილაშვილი....1982: მერქნიანი ჯიშებისა და ზოგიერთ სუბტროპიკულ მცენარეთა ლექსიკონი. გამომცემლობა „განათლება“,

თბილისი.

- 11 . ბათუმის2012 : ბათუმის ბოტანიკური ბაღის ხეები და ბუჩქები
(ანოტირებული სია).ISBN -978-9941-9296-0-1 გამომც.
შ. პ. ს. „საჩინო“, თბილისი. გვ. 111.
12. ბიძინაშვილი.... 2016: ბიძინაშვილი რ. ცხადაძე ნ. ხაიკაშვილი ხ. “
სამკურნალო მცენარეების კოლექცია საქართველოს
ეროვნულ ბოტანიკურ ბაღში“, II სამეცნიერო
კონფერენციის მასალები, ბიომრავალფეროვნება და
საქართველო საქართველოს ეროვნული ბოტანიკური
ბაღი , თბილისი, გვ. 69.
13. ბიძინაშვილი ...2013: ბიძინაშვილი რ. ცხადაძე ნ. ხაიკაშვილი ხ.
თბილისის მიდამოებში მოზარდი ენდემური,
რელიქტური და იშვიათი სამკურნალო მცენარეები/
საქართველოს ეროვნული ბოტანიკური ბაღის შრომები,
თბილისი, გვ. 96.
14. ბათუმის 2012: ბათუმის ბოტანიკური ბაღი 100 წლისაა. „ მცენარეთა
ფიზიოლოგიის და ბიოქიმიის განყოფილება“ მონოგრაფია.
შ პ ს „ საჩინო“. თბილისი. გვ. 49.
15. ბრეგვაძე1974: ბრეგვაძე მ. ბათუმის ბოტანიკურ ბაღში
ინტროდუცირებულ დაფნისებრთა ოჯახის ზოგიერთი
სახეობის ზამთრის სვენები თავისებურებები ბათუმის
ბოტანიკური ბაღის მოამბე N20. გამც. თბილისი, . გვ 82-.
16. ბრეგვაძე....2012: ბრეგვაძე მ.- მეტრეველი მ. იშვიათი ეგზოტის
Cinnamomum japonicum Sieb. Et Nakai ინტროდუქცია,
ადაპტაციის ეკოფიზიოლოგიური ასპექტები და

კონსერვაცია ბათუმის ბოტანიკურ ბაღში. ბათუმის

ბოტანიკური ბაღის მოამბე № 34. თბილისი გვ. 112.

17. ბოლქვაძე....2008: ბოლქვაძე გ. „მარადმწვანე მაღალდეკორატიულ მცენარეთა ბიოეკოლოგიური თავისებურებანი ბათუმის ბოტანიკურ ბაღში“-სადოქტორო დისერტაცია, შოთა რუსთაველის სახელმწიფო უნივერსიტეტი, ბათუმი. გვ. 80 - 89.
18. გოგოძე 1968: გოგოძე ქ. ცინეოლის შემცველობა ევკალიპტის ზოგიერთ სახეობათა ეთეროვან ზეთში, ბათუმის ბოტანიკური ბაღის მოამბე № 13, გამომცემლობა თბილისი, გვ. 94-98.
19. გოგოძე1998: გოგოძე ქ. ეთეროვანი ზეთების შემცველობა ევკალიპტის ზოგიერთი სახეობის ფოთლებში აჭარის პირობებში. ბათუმის ბოტანიკური ბაღის მოამბე, ტ. 30-31. გამომცემლობა „მეცნიერება“ გვ. 40-42.
20. დავითაძე....2002: დავითაძე მ. „აჭარის ადვენტური ფლორის ბიომორფოლოგიური ანალიზი“. გამომცემლობა „ბათუმის უნივერსიტეტი“. ბათუმი, გვ. 13-15.
21. მანჯავიძე..... 1961: მანჯავიძე დ. „ბათუმის მიდამოებში კლიმატური პირობების ზოგიერთი თავისებურებები და მისი გავლენა მერქნიან მცენარეებზე“. ბათუმის ბოტანიკური ბაღის მოამბე №9-10. საქართველოს მეცნიერებათა აკადემიის გამომცემლობა თბილისი. გვ. 175-176.
22. მანჯავიძე...1968: მანჯავიძე დ., ციციძე ა., მატინიანი ა., მოროზოვა გ., თათარიშვილი ა ტატიშვილი ა. „ბათუმის

ბოტანიკური ბაღის ხეები და ბუჩქები“,
ანოტირებული სია, გამომცემლობა „მეცნიერება“.
თბილისი. გვ. 5-153.

23. მაყაშვილი....1961: მაყაშვილი ა. ბოტანიკური ლექსიკონი გამომც. საბჭოთა
საქართველო

24. მირზაშვილი...1933: მირზაშვილი ვ.ი. „მერქნიანი ეგზოტების
აკლიმატიზაცია საქართველოს პარკებში“. თბილისი.

25. საღარეიშვილი... 2008 : თამარ საღარეიშვილი „ფენოლური ნაერთები და
ეთეროვანი ზეთები საქართველოში მოზარდ და
ინტროდუცირებულ ზოგიერთ უმაღლეს მცენარეში“
უაკ 547.972 + 615.9. ISBN 978-9941-0-0607-4, თბილისი,
გ. 29-40.

26. საბაშვილი.....1965: საქართველოს ნიადაგები. თბილისი. გვ.350.

27. სტანდარტი1954: სახელმწიფო სტანდარტი 5055 -49 „თესლის
ხარისხის განსაზღვრის მეთოდები“ საქართველო,
ბათუმის ბოტანიკური ბაღი, გამომც. სახელგამი,
გვ. 42-45.

28. სოხაძე ...2016: სოხაძე ე. ჭანტურია -კვირტია თ. საქართველოში
არსებული მცენარეების-სანელებლების სამკურნალო
თვისებები. ბიომრავალფეროვნება და საქართველო - II
სამეცნიერო კონფერენციის მასალები , საქართველოს
ეროვნული ბოტანიკური ბაღი , თბილისი, გვ. 83.

29. ტყავაძე1990: ტყავაძე მ. კილაძე ი. „ დეკორატიული დენდროლოგია“
II ნაწილი. სოხუმი. გვ. 73.

30. ციცვიძე ...1956: ციცვიძე ა. „ცრუქაფურის ხის ბიოლოგიის
საკითხისათვის“, ბათუმის ბოტანიკური ბაღის მოამბე,

ბათუმი, გვ. 138-140.

31. ციციძე....1958: ციციძე ა. აჭარის შავი ზღვის სანაპიროზე
ინტროდუცირებული ზოგიერთი ეგზოტიკური ჯიშის
აკლიმატიზაციის საკითხებისათვის, ბათუმის
ბოტანიკური ბაღის მოამბე №8, თბილისი. გვ. 179-191.
- 32 . ციციძე ... 1960: ციციძე ა., ჰიმაღლის მცენარეთა ინტროდუქცია-
აკლიმატიზაციის შედეგები აჭარაში.“ ბათუმის
ბოტანიკური ბაღი, მოამბე № 9-10, გვ. 125-127.
33. ჩაიძე...2009: ჩაიძე ფ. „ აჭარის ზღვისპირეთში გავრცელებული გვარი *Magnolia*
L ინტროდუქციის შედეგები“. სადოქტორო დისერტაცია, შოთა
რუსთაველის სახელმწიფო უნივერსიტეტი, ბათუმი. 40-70.
34. ჩაიძე....2016: ჩაიძე ფ. აბაშიძე ნ. ქათამაძე თ, აღმოსავლეთაზიური
წარმოშობის სამკურნალო მცენარეთა გენოფონდი
ბათუმის ბოტანიკურ ბაღში. ბიომრავალფეროვნება და
საქართველო - II სამეცნიერო კონფერენციის მასალები ,
საქართველოს ეროვნული ბოტანიკური ბაღი ,
თბილისი, გვ. 80-82.
35. ჩხაიძე.... 1996: ჩხაიძე გ. „ სუბტროპიკული კულტურები“, თბილისი,
გვ. 4-20.
36. ჯაბნძე...2004: ჯაბნძე რ. „ჩაი და ციტრუსები.“ თბილისი. გვ. 99-104.
37. Абашидзе ..2015 Абашидзе Н. Г., Каландия А.Г, Чаидзе Ф. Э.
„Биоэкологические и био-химические особенности
лептоспермумов (*Leptospermum* J.R.Forst. & G.Forst.)
интродуцированных в Батумском ботаническом саду.“
Материалы III-й Международной научно-практической
конференции «Проблемы сохранения биологического
разнообразия и использования биологических ресурсов»,

посвященной 110-летию со дня рождения академика
Н. В.Смольского. 7–9 октября 2015 г. Минск. Беларусь. В
двух частях. Часть I. Минск «Конфидо» ISBN 978-985-
6777-74-8. С. 253-255.

38. Академия....1972: Академия наук СССР, Главный ботанический сад,
„Интродукционные фонды юго-восточноази“. Изд.
„Наука“, Москва, стр. 188-198.
39. Алаев....1957: Алаев Б.С. „Эфирномасличные растения и парфюмерно-
косметические предприятия Китая“. Москва, стр. 56.
40. Алексеев...1955 : Алксеев В. П. Растительные ресурсы Китая.
Издательство всесоюзного растениеводства.
Ленинград, стр. 66.
41. Алексеева1953 : Алексеева Е. А. „Эфирномасличные культуры южной
полосы СССР". Государственное издательство
сельскохозяйственной литературы. Москва. ст. 14-40.
42. Анненкова1961: Анненкова Г. А. "Сборник научно-исследовательских
работ по масличным и эфиромасличным культурам".
Издательство Советская Кубань. Ст. 142-216.
43. Алехин1944: Алехин В. В. „География растений“. Москва, стр. 22.
44. Аннотированный....1987: Аннотированный список «Деревьяикустарники
БатумскогоБотанического сада», издательство
«Мецниереба», стр. 23-63.
45. Атлас1962 : Атлас лекарственных растений СССР. Москва.
»Наука».
46. Батумский2007: Батумский Ботанический сад. «Деревья и
кустарники Батумского Ботанического сада
(Покрытосемянные)“ . Часть II, стр. 54-80.
47. Бейдеман...1974: Бейдеман И.Н. "Методика изучения фенологии

растений и растительных сообществ", изд. "Наука",
Новосибирск, стр. 78-90.

48. Био-химические ...1972 Био-химические методы анализа эфиромасличных
растений Всесоюзный научно-исследовательский институт
эфиромасличных культур. Симферополь. 1972. Ст. 194-196.
49. Брегвадзе ... 1984: Брегвадзе М. А. Шарашидзе Н. М. Характер ростовых
процессов и приспособления растений к новым
условиям среды. Тезиси докладов Всесою.Совещан.
»Актуальные задачи физиологии и биохимии растений
в ботанических садах СССР. » Пушкино.с.24.
50. Буйко1968 : Буйко Р. А., Маковкина А. И., Соколов В. С.
Библиография по эфирномасличным растениям и
эфирочным маслам –указатели отечественной
литературы за 1747-1965 гг. Издательство "Наука".
Ленинград. .
51. Гурский.....1957: Гурский А. В. Основные итоги интродукции древесных
растений в СССР. Изд-во АН. СССР М. Ленинград. С. 303.
52. Дараселия 1949: Дараселия М.К. „Красноземные и подзолистые почвы
Грузии и их использования под субтропические
культуры“. Махарадзе-Анасеули.
53. Дворецкий1964: Дворецкий М.Л. “Текущий прирост древесного ствола и
древостоя. Из. “Лесная промышленность“. Москва.
54. Доспехов.....1985 : Доспехов В.”Методика полевого опыта”. Москва,
Агропромиздат, стр. 351.
55. Елагин.....1979: Елагин И. Н. Лобанов А. И. Атлас определитель
фенологических фаз растений. М. изд. »Наука». С. 95.
56. Вассерман1939: Вассерман И. Г. “ Интродукция эфирномасличных

растений." Издательство " Виэмп." Пушкино-
Московский обл, стр. 10-12.

57. Вульф..... 1933: Вульф Е. В. В.И.Нилов, " Эфирномасличные растения их
культура и эфирные масла . " Том I. Издания
всесоюзного института растениеводства. Ленинград.
58. Вульф..... 1934: Вульф Е. В., Нилов В.И. " Эфирномасличные растения .с. 50.
их культура и эфирные масла . " Том II. Издания
всесоюзного института растениеводства. Ленинград. С.43.
59. Вульф..... 1937: Вульф Е. В. В.И.Нилов, " Эфирномасличные растения .
их культура и эфирные масла . " Том III. Издания
всесоюзного института растениеводства. Ленинград. С.31.
60. Гераськин..... 2010: Гераськин С.А., Сарануцева Е.И., Цаценко А.В. и др.
„Биологический контроль окружающей среды“.
„Генетический мониторинг“, Москва. С. 45-49.
61. Гинкул.....1936: Гинкул С. Г., „Интродукция и акклиматизация растений
влажных субтропиках СССР.“ Известия Батумского
субтропического сада. №1. Батуми. С. 3-44.
62. Гинкул..... 1940: Гинкул С. “Итоги интродукция растений в Батумском
Ботаническом Саду”. Изв. Батумского Субтроп. Ботанического
сада, № 5. Батуми. С. 3-51.
63. Головкин....1981: Головкин Б. Н. „История интродукции растений в
Ботанических садах,, издательство московского
Университета, Москва. С. 7.

64. Гост 1979 : Гост 14618-12-78. "Масла эфирные, вещества душистые и полупродукты их синтеза." Гост Гост 14618-12-78.
Москва, ст. 103.
65. Дмитриева1947: Дмитриева А. А. Путеводитель по Батумского Субтропическому ботаническому саду. Батуми. С. 3-56.
66. Иванова.....1982: Иванова З. Я. „Биологические основы и приемы вегетативного размножения древесных растений стеблевыми и черенками.» Киев. Академия наук украинской ССР, Центральный республиканский ботанический центр. С. 34-38.
67. Карманова....1976: Карманова И. В. – Математические методы изучения роста и продуктивности растений- Академия наук СССР- Лаборатория лесоведения ю Изд. Наука. Москва.с. 25-29.
68. Краснов..... 1913: Краснов А. Н. „ Батумский ботанический сад за первый год организации», Журнал. Русские субтропики. Батуми. №9. 3-7.
69. Красюк1929: Красюк А. „ Почвы и их исследование в природе.“ Государственное издательство, с. 20.
70. Крокер....1955: Крокер В., Бартон Л., "Физиология семян". Изд. иностранной литературы, Москва, стр. 20-26.
71. Колесниченко...1976: Колесниченко М. В. -“Биохимические взаимовлияние древесных растений“. Изд. “Лесная промышленность.“ Москва.
72. Кохреидзе1938: Кохреидзе В. Г. „Фенология субтропических растений“, Издание Батумского субтропического Ботанического сада. Батуми, с. 2-43.
73. Кудинов1984: Кудинов М., Кухарева Л. „Новые эфирномасличные растения в Белорусии. Академия наук БССР центральный ботанический сад. Минск. Ст. 45-58.
74. Лапин.....1984: Лапин В. И. Академия наук СССР. Главный

ботанический сад. „Роль Интродукции в сохранении генофонда редких исчезающих видов растений.“
Изд. “ Наука“. Москва. С.4.

75. Лукашова1958: Лукашова Е. Н. „Южная Америка“. Государственная учебно педагогическая издательство министерства просвещения РСФСР. Москва.13.
76. Методика.....1975: Методика фенологических наблюдении в ботанических садах СССР. С. 27.
77. Минкевич1949: Минкевич И. Берковский В. Е. "Масличные культуры". Государственное издательство сельскохозяйственной литературы. Москва, Ст. 32-154.
78. Миронова.....1984: Миронова Ю. С. „ Кудrania трехлопастная (Cudrania tricuspidata (Carr.) Burr.) в Ботаническом саду АН УзССР. . ISSN 0135-1664. Вып. 19. Изд. „ Фан„ Узбекской ССР, Ташкент. С. 83.
79. Мичурин...1936: Мичурин И. В. „ Роль акклиматизации в распространение растений“. Москва. изд. наука. С. 39-44.
80. Мичурин.....1955: Мичурин И. В. „Акклиматизация растений“. Москва. Сельхозгиз. С.32.
81. Молчанов...1967: Молчанов А. А. "География плодоношения главнейших древесных пород в СССР“. Академия наук СССР. Лаборатория лесоведения. Изд. “Наука“. Москва. с. 11-99.
82. Молот ...1938: Молот К. Г. Труды интродукционного питомника субтропических культур. Выпуск 9. " Камфорное дерево". Ст. 43.
83. Морозова1957: Морозова Г. А. " Ритм роста восточноазиатских растений

в Батумском ботаническом саду. »Известия» №8

Батумского Ботанического сада, Издательство Академия наук Грузинской ССР, Батуми, стр. 34-50.

84. Морозова1968: Морозова Г. «Испытание новых видов Восточноазиатских растений» Издательство «Мецниереба» Тбилиси, № 13. С. 30-40.
85. Некрасов..... 1980: Некрасов В. И. „Актуальные вопросы развития теории акклиматизации растений“. Изд. " Наука". Москва. Стр. 10-11.
86. Нестеровичь...1950: Нестеровичь Н. Д. „ О проращивании пыльцы древесных пород в связи с их плодоношением“ . Изв. АН БССР. С. 150-130.
87. Нестеровичь.... 1969: Нестеровичь Н. Д., Маргаилик Г.И. “Влияние света на древесные растения“. Изд. “Наука и техника“. Академия наук СССР- институт экспериментальной ботаники. С. 46-80.
88. Нестеровичь.... 1972: Нестеровичь Н. Д. „Древесные растения и влажность почвы“. Изд. „ Наука и техника“, Минск. С. 25-67.
89. Нижарадзе 1961: Нижарадзе Н. „Советская Аджария. Экономико-Географическая характеристика“. Батуми, с. 254.
90. Папунидзе..... 1987: Папунидзе В. Кипнис М. „Почвенный покров Батумского Ботанического сада“. Изв . Ббс. Тбилиси: «Мецниереба» Тбилиси, ст. 228.
91. Петров.....1978: Петров И. П. „Интродукция древесных растений Средней Азии в Москве „. Издательство «Наука».
92. Пилипенко.... 1978: Пилипенко ф. «Иноземные деревья и кустарники на Черноморском побережье Кавказа“. Издательство «Наука». С. 293.
93. Прянишников1934: Прянишников Д.Н. Агрохимия. (Учебник). М.; Л.: Сельхозиздат, 399 с.

94. Протопопов....1973: Протопопов Г, Ф., Ткаченко В, И., Купченко А. И.
°Созревание семян древесных и кустарниковых
растений, Издательство ° Наука°, Фрунзе , ст. 13-67.
95. Раскатов.....1974: Раскатов П. Б. "Анатомия вегетативных органов древесных
растений". Изд. Воронежского университета. Воронеж.
96. Рутовский....1925: Рутовский Б. Н., Кутовский Б. Н. «Душистые растения и
эфирные масла Крыма и Кавказа ». Труды Научного Химико-
фармацевтического института. Выпуск II Москва. Ст. 50-117.
97. Сабашвили 1968: Сабашвили М. Н. „Почвы Грузии“. Тбилиси. Агротехника
масличных культур. Сборник научных работ. Отдел
земледелия. Краснодар. Ст. 26-46.
98. Словарь...1984 : Словарь географических названий СССР. Издание второе,
переработанное и дополненное. Москва."Недра". с. 13-37.
99. Смолянова....1976: Смолянова А. М. „Эфирномасличные культкры". Москва.
Издательство "Колос". 1976. Ст. 301-328.
100. Серебряков....1952: Серебряков И.Г. Морфология вегетативных органов
высших растений. Москва С. 388.
101. Серебряков....1953: Проблемы морфологии покрытосемянных. Автореф. док. дисс.
М. 32-76.
102. Серебряков....1954: Серебряков И.Г. О методах изучения ритмики сезонного
развития растений в геоботанических стационарах. Уч. зап.
МГПИ им. В. П.Потомкина, т.37, каф. Биология, вып.2. С. 5-22.
103. Серебряков....1959: Типы развития побегов у травянистых многолетников и
факторы их формирования. Уч. зап. МГПИ им. В. П.
Потомкина, каф. бот.вып. 5. С. 43-65.

104. Серебряков....1964: Сравнительный анализ некоторых признаков ритма
сезонного развития растений различных ботанико-
географических зон СССР. Бюлл. МОИП, отд.
биол.т.69, вып. 5. С. 5-43.
105. Серебряков....1968: Побегообразования и жизненные формы некоторых овсяниц
(Festsuca L.) в связи их эволюцией. В сб. „ Вопросы морфогенеза
цветковых растений и строение их популяций“. Изд. „Наука“,
М. 76-87.
106. Серебряков... 1971: Морфогенез побегов и эволюция жизненных форм злаков.
Изд. „Наука“; М. С. 120-165.
107. Соколов ...1952: Соколов В. С. Алкалоидоносные растения СССР. Изд.
Академия наук СССР. Изд. Ленинград. Стр. 106-110.
108. Соколов...1956: Соколов Д. °Состав плодов мыльного дерева°, Известия
Батумского Ботанического сада, Батуми , ст. 117-120.
109. Супрунов ...1972: Супрунов Н. Горовой Н. Ю. Панков, “Эфирно-масличные
растения дальнего востока” Изд. “Наука”, Новосибирск. 20-77.
110. Тахтаджян ...1966: Система и филогения цветковых растений. М. Л. Изд. «Наука».
111. Тахтаджян1978: Флористические области земли Л. .» Наука». С. 246.
112. Ткачук.....2004: Ткачук Л. П. „Онтогенез *Psidium littorale Raddif*“
В условиях Защищенного грунта“. ISSN 1728-6204.
Промышленная ботаника, Вып 4, стр. 3-8.
113. Ткаченко....2008 : Ткаченко К. Г., Казаринова Н. В. " Использование растений
для санации помещений и профилактики инфекционных
заболеваний. Научные ведомости N 6(46) УДК 581.6:58. 072

613:614:616-093 /098. ст. 82.

114. Тимирязев....1948: Тимирязев К. А. Земледелие и физиология растений ОГИЗ - СЕЛЬХОЗГИЗ, М., 1948 г. Избранные сочинения в 4-х томах.
115. Тупик....2014: Тупик П. В. Интродукция древесных видов, Курс лекций для студентов специальности «Лесовосстановление и питомническое хозяйство» . УДК 581.1(075.8) ББК 43.4я73 Т85. УО «Белорусский государственный технологический университет». Белорус. 3-12.
116. Шарапов.... 1959: Шарапов Н. И. „Масличные растения и маслообразовательный процесс». Издат. академия наук СССР – Москва- Ленинград, Ст. 275 -385.
117. Шарашидзе ... 1985: Шарашидзе Н. “ Физиологические особенности адаптации интродукционных растений .“ Батуми, с. 56.
118. Шевченко 2009: Шевченко С.В. „Репродуктивная биология декоративных и субтропических плодовых растений Крыма“. Украинская Академия Аграрных наук. Никитский ботанический сад. Национальный научный центр. ISBN 978-966-540-286-2.93. 12-34.
119. Шульман 1955: Шульман Н, К. „ Природа Новой зеландии. География“, Из. “Наука . ” Москва, стр. 27-25.
120. Усенко....1984: Усенко Н. В. Деревья, кустарники и лианы дальнего востока. Справочная книга. Хабаровское книжное издательство, Хабаровск. Ст. 29-147.
121. Урбах ...1975: Урбах В. „Статистический анализ в биологических и медицинских исследованиях“ Москва, Издательство «Медицина» , стр. 153.
122. Хромова....1980: Хромова Т. В. “ Методические указание по размножению

интродуцированных древесных растений черенками“. Академия наук ССР- Главный ботанический сад. С. 4-29.

123. Чаидзе.... 2015: Чаидзе Ф.Э., Метревели М.В., Абашидзе Н.Г. “Интродукция и *ex-situ* консервация полезных нетрадиционных растений в Батумском ботаническом саду“. Материалы II

Международной научной конференции „Агробιοразнообразие для улучшения питания“

124. Чиков1981: Чиков П. С., Павлов М. С - Наука и лекарственные растения. Изд. „Знание „, Москва.. Ст. 72-81.

125. Холодко.....1976: Холодко В. С. Д. А. Глоба-Миххайленко. «Ценные древесные породы черноморского побережья Кавказа“. Издател. «Лесная промышленность, Москва. 3-54.

126. Цицин1972: Цицин Н. В. « Интродукционные фонды юго- восточной азии». Издат/ “Наука .” Москва, стр, 91-109.

127. Цицин ... 1962: Цицин Н. В.. Атлас лекарственных растений СССР. Государственное издательство медицинской литературы. Москва. Ст. 68-650.

128. Abashidze....2017 N. Abashidze, F. Chaidze, A. Kalandia. Bioecological and content of essential oils of some plants Introduced at Batumi Botanical Garden. International Scientific Conference 29 september-1 octomber Future Technologies and Quality of Life. BSU, BAU- International university Batumi, (Poster).

129. Abashidze ... 2020: Abashidze N.G.1, Chaidze F. E.2, 1 Kalandia A.3 Chikovani D.4 Growth—development and Biochemical Peculiarities of some plants introduced at Batumi Botanical Garden. International

Journal of Environmental Sciences (ISSN: 2277-1948), 47-49.

130. Amélia2017: Amélia C. Tuler^{1 3} Tatiana T. Carrijo² Márcia F.S. Ferreria² Ariane L. Peixoto¹ Flora of Espírito Santo: *Psidium* (Myrtaceae)
Print version ISSN 0370- 6583 *On-line version* ISSN 2175-786
Rodriguésia vol.68 no.5 Rio de Janeiro. International
Trade Centre (ITC) Sustainable Sourcing: Markets for
Certified Chinese Medicinal and Aromatic Plants. Geneva: ITC,
2016. xvi, 141 pages (Technical paper) Doc.
No. SC-2016-5.E
131. Begum....2004: Sabira Begum, Syed Imran Hassan, Syed Nawazishali and
BinS. iddiqui. Chemical constituents from the leaves of
Psidium guahava. H.E.J. Research Institute of Chemistry,
International Center for Chemical Sciences, University of
Karachi, Karachi- 5270, Pakistan.
132. Chaidze ...2015 : Chaidze F. E. Metreveli M. V. Abashidze N. G. Introduction and
Ex-situ Preservation of Useful Non-traditional Plants in the
Batumi Botanical Garden . Slovak University of Agriculture in
Nitra. Agrobiodiversity for improving nutrition, health and life
quality. Part 1. The Scientific proceedings of the international
network Agrobionet . Nitra p 82-84.
133. Dawson 1997: A history of *Leptospermum scoparium* in cultivation –
discoveries from the wild. The New Plantsman 4: 51–59.
134. Denis2014 Denis Wilson, Allan Stiles. “Robertson Rainforest Plant Species
Sassafras (Doryphora sassafras). ISSN 1037 – 2032, Robertson
Environment Protection Society – to promote the protection and
enhancement of the Robertson environment.
PO Box 3045, Robertson NSW 2577. p. 3-6.
135. Douglas....2001: www. „Defining the North Island manuka chemotype

- resource a survey report M Douglas, R Anderson, J van Klink, N Perry B Smallfield, July 2001. Crop & Food Research Report No. 447 New Zealand Institute for Crop & Food Research Limited. P. 5-10. J Agric Food Chem, P. 4-7.
136. Dvorakovskaya...2015: Dvorakovskaya V. M. “ Wild food plants of the far east in the main botanical garden of the ras” Agrobiodiversity for improving nutrition, health and life quality. Part 1. The Scientific proceedings of the international network Agrobionet . Nitra . Ст. 139-144.
137. Garanovich....2015: Garanovich L. M., Shpitainaia T. V. “Cultures of medical horticulture in Belarus.” Agrobiodiversity for improving nutrition, health and life quality. Part 1. Agrobiodiversity for improving nutrition, health and life quality”. Nitra . P. 165-168.
138. Cockayne1958: Cockayne, L. „The vegetation of new Zealand“. Robert A. DeFilipps¹, Gary A. Krupnick². The medicinal plants of Myanmar. Department of Botany, National Museum of Natural History, Smithsonian Institution, Box 37012, MRC- 166, Washington, DC, 20013-7012, USA.
139. Govaerts.....1996: www. Govaerts R. „World checklist and bibliography of Magnoliaceae“. Richmond Royal Botanic Gardens. Kew.
140. Johnson.....1980: The leaf anatomy of *Leptospermum* Forst. (Myrtaceae). Australian Journal of Botany 28: 77–104.
141. José2014: José M. Alvarez-Suarez¹, Massimiliano Gasparrini, Tamara Y. Forbes-Hernández, Luca Mazzoni and Francesca Giampieri The Composition and Biological Activity of Honey: A Focus on Manuka Honey.

142. Harkenthal..1999: www. Harkenthal, M., Reichling, J., Geiss, H. –K.. and Saller, Comparative study on the in vitro antibacterial activity of Australian tea tree oil, cajuput oil, niaouli oil, Manuka oil, kanuka oil, and eucalyptus oil. Pharmazie 54, 460-463
143. Hudz 2015: Hudz N. L, Koretska A. M. Kalyniuk T. G., Korytiuk R. S. „Features of the Pharmaceutical Development of Herbal Medicinal Products Agrobiodiversity for improving nutrition, health and life quality. Part 1. Agrobiodiversity for improving nutrition, health and life quality”. Nitra. P. 262-267.
144. Katyayani2012 Katyayani Dutta Choudhury¹, M. Dutta Choudhury², S. B. Paul¹ & M. K. Baruah³ „Bioactivities of some ethno-medicinal Rubiaceae plants available from Assam,, – a review. 1 Ethno- botany & Medicinal plant research laboratory, Department of Life Science, Assam University. Silchar -7880011, Assam, India. 2 Department of Chemistry, Assam University. Silchar – 7880011, Assam, India. 3 Department of Botany, Cachar College, Silchar 788001, Assam, India.
145. Kuklina....2015: Kuklina A. G. Vinogradova Yu. K., Shelepova O.V. “The Possibility of invasive Plant ’s Application for Medicinal Purp. Agrobiodiversity for improving nutrition, health and life quality. Part 1. Agrobiodiversity for improving nutrition, health and life quality” Nitra . P. 399-401.
146. Mayr..... 1909 : Mayr H. Die Naturgesetzlcher Grundlage des Waldbaues. Berlin: Parey, 366 S.
147. Molecules2013 : Francisco J. Luna vizquez, Cesar Ibarra-Alvarado, Alejandra Rojas-Molina, Juana I. Rojas-Molina, „ Nutraceutical Value of Black Cherry Prunus serotina

- Ehrh. Fruits:Antioxidant and Antihypertensive Properties“, *Molecules* ISSN 1420-3049.
148. Perry....1997: www. Perry. N. B., Brennan, N. J., van Klink, J. W. Harris, W., Douglas, M. H. McGimpsey, J.A. Smallfield, B.M. and Anderson, R. E. Essential oils from New Zealand Manuka and kanuka: Chemotaxonomy of *Leptospermum*. *Phytochemistry* 44, 1485-1494.
 149. Porter,.....1998: www. Porter, N.G. and Wilkins, A. L. Chemical, physical and antimicrobial properties of essential oils of *Leptospermum scoparium* and *Kunzea ericoides*. *Phytochemistry* 50, 407-415.
 150. Porter.... 1998 N. G. Porter a , P. E. Smale b , M. A. Nelson b , A. J. Hay a , J. W. Van Klink c & C. M. Dean- Variability in essential oil chemistry and plant morphology within a *Leptospermum scoparium* population New Zealand Journal of Botany, Vol. 36: 125-1330028-825X/98/3601-125. The Royal Society of New Zealand.
 151. Richard....1902: Richard T., Baker F. “ A research on the Eucalyptus, especially in regard on their Essential oils.
 152. Stephens....2006 : The factors responsible for the varying levels of UMF® in mānuka (*Leptospermum scoparium*) honey. University of Waikato Hamilton, New Zealand. P. 12-15.
 153. Tameike ...1895 : Tameike, Akasaka. “Useful plants of Japan”. Agricultural society of Japan Tokio. 3-43.
 154. Thompson.... 1989: A revision of the genus *Leptospermum*. *Telopea* 3: 301–448.
 155. William ...2016: William N Setzer Chemical composition of the leaf essential oil of *Lindera benzoin* growing in North Alabama Department of Chemistry, University of Alabama in

- Huntsville Huntsville, AL 35899, USA . . American journal of essential oils and natural products, ISSN: 2321 9114.
156. Wilker1999: Wilker C., Pedrosa-Macedo J. H., Vitorino M. D., Caxambu M. G., Smit C. W. h, Decie - Unicentro – „ Strawberry Guava (*Psidium cattleianum*) – Prospects for Biological Control, “ Universidade do Centro-Oeste, PR 153 - KM 7 - Bairro Riozinho - C. Postal 21, 84500-000 - Irati – PR Brazil. Proceedings of the Symposium on biological Control of Weeds 659, 4-14 July 1999, Montana State University, Bozeman, Montana, USA Neal R. Spencer . pp. 659-665.
157. Yamaguchi ...1982: www. Yamaguchi K, Shibamoto T. Isolation and identification of banana-like aroma from banan shrub (Micheli figo). J Agric Food Chem, P. 4-749.
158. Yang.....2017: Yang D, et al. www.hong Yao Cai.
159. <https://www.IUCNredlist.org>.
160. <https://www.theplantlist.org>.